

**Access Eligibility Criteria for Infants and Toddlers Who Require Specialty Services and Supports
(birth through 3 years [47 months] of age)**

Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+

March 18, 2008

DC 0-3 R Diagnosis	ICD 9 CM Diagnosis <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>	DSM IVR Diagnosis
100 Post traumatic Stress Disorder	<p>308 Acute reaction to stress** (requires 4th digit subclassification listed below). Includes:</p> <ul style="list-style-type: none"> • Catastrophic stress • Gross stress reaction (acute) • Transient disorders in response to exceptional physical or mental stress which usually subside within hours or days <p>308.0 Predominant disturbance of emotions</p> <ul style="list-style-type: none"> • Anxiety as acute reaction to exceptional (gross) stress <p>308.1 Predominant disturbance of consciousness Fugues as acute reaction to exceptional (gross) stress</p> <p>308.2 Predominant psychomotor disturbance</p> <ul style="list-style-type: none"> • Agitation states as acute reaction to exceptional (gross) stress • Stupor as acute reaction to exceptional (gross) stress <p>308.3 Other acute reactions to stress**</p> <ul style="list-style-type: none"> • Acute situational disturbance • Brief or acute posttraumatic stress disorder <p>308.4 Mixed disorders as reaction to stress</p> <p>308.9 Unspecified acute reaction to stress</p> <p>309.81 Prolonged posttraumatic stress disorder**</p> <ul style="list-style-type: none"> • Chronic posttraumatic stress disorder 	<p>Acute Stress Disorder</p> <p>308.3</p> <p>Post traumatic Stress Disorder</p> <p>309.81</p>
150 Deprivation/Maltreatment Disorder	<p>313.89 Emotional Disturbance of Childhood NEC**</p> <p>313.9 Unspecified emotional Disturbance of Childhood or adolescence</p>	<p>313.89 Reactive Attachment Disorder of Infancy or Early Childhood</p>

**Access Eligibility Criteria for Infants and Toddlers Who Require Specialty Services and Supports
(birth through 3 years [47 months] of age)**

Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+

March 18, 2008

DC 0-3 R Diagnosis		ICD 9 CM Diagnosis <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>		DSM IVR Diagnosis	
200	Disorder of Affect				
210	Prolonged Bereavement/Grief Reaction	309.00	Adjustment disorder with depressed mood • Grief reaction	309.0	Adjustment Disorder with Depressed Mood
		309.1	Prolonged depressive reaction**		
220	Anxiety Disorder of Infancy & Early Children	300.00	Anxiety state, unspecified** • Anxiety: neurosis, reaction, state (neurotic) • Atypical anxiety disorder	300.00	Anxiety Disorder NOS
		300.01	Panic disorder. Panic: attack, state	300.01	Panic disorder without
		300.02	Generalized anxiety disorder	300.02	Generalized Anxiety Disorder
		309.21	Separation anxiety disorder	309.21	Separation Anxiety Disorder
		313.0	Overanxious disorder** • Anxiety and fearfulness of childhood and adolescence • Overanxious disorder of childhood and adolescence		
221	Separation Anxiety	309.21	Separation anxiety disorder	309.21	Separation Anxiety Disorder
222	Specific Phobia	300.29	Other isolated or specific phobias	300.29	Specific Phobia
223	Social Anxiety Disorder	300.23	Social Phobia	300.02	Social Phobia
		313.2	Sensitivity, shyness and social withdrawn disorder (requires 5 th digit subclassification—listed below)		
		313.21	Shyness disorder of childhood		
		313.22	Sensitivity reaction of childhood or adolescence Introverted disorder of childhood --Social withdrawal of childhood or adolescence --Withdrawal reaction of childhood or adolescence		
224	Generalized Anxiety Disorder			300.02	Generalized Anxiety Disorder
225	Anxiety Disorder NOS			300.0	Anxiety Disorder NOS

Access Eligibility Criteria for Infants and Toddlers Who Require Specialty Services and Supports
(birth through 3 years [47 months] of age)

Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+

March 18, 2008

DC 0-3 R Diagnosis		ICD 9 CM Diagnosis <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>		DSM IVR Diagnosis
230	Depression of Infancy and Early Childhood			
231	Type I Major Depression	296.2X Major Depressive Disorder, Single Episode, Unspecified 5 th digit: 0 unspecified 1 mild, 2 moderate, 3 severe without mention of psychotic behavior, 4 severe with specified as with psychotic behavior, 5 in partial or unspecified remission, 6 in full remission	296.20	Major Depressive Disorder, Single Episode, Unspecified
232	Type II Depressive Disorder NOS	296.3X Major Depressive Disorder, Recurrent, Unspecified 5 th digit: same as above	296.30	Major Depressive Disorder, Recurrent, Unspecified
		311 Depressive Disorder, not elsewhere classified --Depressive Disorder NOS --Depressive State NOS--Depression NOS	311	Depressive Disorder NOS
		313.1 Misery and unhappiness disorder		
240	Mixed Disorder of Emotional Expressiveness	300.4 Neurotic Depression --Anxiety depression --Depression with anxiety --Depressive reaction --Dysthymic Disorder --Neurotic depressive State --Reactive Depression	300.4	Dysthymic Disorder
		313.9 Unspecified emotional disturbances of childhood or adolescence	313.9	Disorder of Infancy, Childhood, or Adolescence NOS

**Access Eligibility Criteria for Infants and Toddlers Who Require Specialty Services and Supports
(birth through 3 years [47 months] of age)**

**Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+
March 18, 2008**

DC 0-3 R Diagnosis	ICD 9 CM Diagnosis <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>		DSM IVR Diagnosis
300 Adjustment Disorder	309 309.0	Adjustment Reaction (309.0—309.9) Brief depressive reaction --Adjustment disorder with depressed mood --Grief reaction	309.9 Adjustment Disorder: Unspecified
	309.24	Adjustment reaction with anxious mood	309.24 Adjustment Disorder with Anxiety
	309.28	Adjustment reaction with mixed emotional features	309.28 Adjustment Disorder with Mixed Anxiety and Depressed Mood
	309.3	With predominant disturbance of conduct --conduct disturbance as adjustment reaction --destructiveness as adjustment reaction	309.3 Adjustment Disorder with Disturbance of Conduct
	309.4	With mixed disturbances of emotions and conduct	309.4 Adjustment Disorder with Mixed Disturbance of Emotions and Conduct
	309.29	Other --culture shock	309.9 Adjustment Disorder NOS
	309.82	Adjustment reaction with physical symptoms	309.9 Adjustment Disorder NOS
	309.83	Adjustment reaction with withdrawal --Elective mutism as adjustment reaction	313.23 Selective Mutism
	309.9	--Hospitalism (in children) NOS Unspecified adjustment reaction --Adaptation reaction NOS --Adjustment reaction NOS	309.9 Adjustment Disorder NOS 309.9 Adjustment Disorder NOS
400* – Regulation Disorders of Sensory Processing Sensory Processing when coupled with parent/child interaction	312.9 313.0	Unspecified disturbance of conduct Overanxious disorder** --Anxiety and fearfulness of childhood and adolescence --Overanxious disorder of childhood and adolescence	312.9 300.02 Disruptive Behavior Disorder NOS Generalized Anxiety Disorder
410* Hypersensitive			
411* Type A: Fearful/Cautious			

+ To be used as a guide in Clinical Assessment/Diagnosis

Access Eligibility Criteria for Infants and Toddlers Who Require Specialty Services and Supports
(birth through 3 years [47 months] of age)

Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+

March 18, 2008

DC 0-3 R Diagnosis	ICD 9 CM Diagnosis <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>	DSM IVR Diagnosis
412*	Type B: Negative/Defiant 312.81 312.9 313.81 314.00	Conduct Disorder—Child onset (Note: for young child over 36 months of age) Disruptive Behavior Disorder NOS Oppositional Defiant Disorder Attention Deficit Disorder— Predominantly inattentive type
420*	Hypo/Underresponsive 314.0 314.01	Attention Deficit Disorder --Requires 5 th digit subclassification--listed below With hyperactivity--combined type --Overactivity NOS --Predominantly hyperactive/impulsive
430*	Sensory Stimulation – Seeking/Impulsive 314.9	Impulse Control Disorder NOS ADHD-Hyperactive impulsive disorder Attention Deficit/Hyperactivity Disorder NOS
500*	Sleep Disorder	
510*	Sleep Onset Disorder 307.40 307.41	Dyssomnia NOS or Parasomnia NOS Insomnia Related to....
	Nonorganic sleep disorder, unspecified	307.47
	Transient disorder of initiating or maintaining sleep • Hyposomnia associated with acute or intermittent emotional reactions or conflicts • Insomnia associated with acute or intermittent emotional reactions or conflicts • Sleeplessness associated with	327.02

**Access Eligibility Criteria for Infants and Toddlers Who Require Specialty Services and Supports
(birth through 3 years [47 months] of age)**

**Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+
March 18, 2008**

DC 0-3 R Diagnosis	ICD 9 CM Diagnosis <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>	DSM IVR Diagnosis
	307.42 Persistent disorder of initiating or maintaining wakefulness • Hyposomnia, insomnia, or sleeplessness associated with: anxiety, conditioned arousal, depression (major) (minor), psychosis	307.42 or 327.02 Primary Insomnia Or Insomnia Related to...
	307.43 Transient disorder of initiating or maintaining wakefulness • Hypersomnia associated with acute or intermittent emotional reactions or conflicts	307.15 Hypersomnia Related to....
	307.44 Persistent disorder of initiating or maintaining wakefulness • Hypersomnia associated with depression (major) (minor)	307.44 or 327.15 Primary Hpersomnia or Hypersomnia Related to....
520*	307.45 Circadian Rhythm Sleep Disorder of nonorganic origin	307.30 or 327.31 Circadian Rhythm Sleep Disorder, Unspecified type or Delayed Sleep Phase Type
	307.46 Somnambulism or night terrors	307.46 Sleep Terror Disorder or Sleepwalking Disorder
	307.47 Other dysfunctions of sleep stages or arousal from sleep Nightmares: NOS, REM-sleep type, sleep drunkenness	307.47 Dyssomnia NOS, Nightmare Disorder, or Parasomnia NOS
	307.48 Repetitive intrusions of sleep Repetitive intrusions of sleep with: atypical polysomnographic features, environmental disturbances, repeated REM-sleep interruptions	307.47 Parasomnia NOS
	307.49 Other "Short sleeper," subjective insomnia complaint	307.42 Primary Insomnia
600	Feeding Behavior Disorders	
601	Feeding Behavior Disorders Other • Infantile feeding disturbances of	307.59 Feeding Disorder of Infancy or Early Childhood

+ To be used as a guide in Clinical Assessment/Diagnosis

Access Eligibility Criteria for Infants and Toddlers Who Require Specialty Services and Supports
(birth through 3 years [47 months] of age)

Crosswalk between Diagnostic Classifications 0-3, ICD 9 CM and DSM IVR+

March 18, 2008

DC 0-3 R Diagnosis	ICD 9 CM Diagnosis <i>** Note: Refer to ICD 9 Coding Manual for Exclusions to this Diagnosis</i>	DSM IVR Diagnosis
602 Feeding Disorders of Caregivers Caregiver/Infant Reciprocity	nonorganic origin • Loss of appetite—of nonorganic origin Other • Infantile feeding disturbances of nonorganic origin • Loss of appetite—of nonorganic origin	307.59 Feeding Disorder of Infancy or Early Childhood
603 Infantile Anorexia	307.59 Other • Infantile feeding disturbances of nonorganic origin • Loss of appetite—of nonorganic origin	307.59 Feeding Disorder of Infancy or Early Childhood
800 Other Disorders		

* Not Axis I Diagnoses

References

American Psychiatric Association, Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR) Fourth Edition (Text Revision), 2000.

National Center for Health Statistics, International Classification of Diseases, Ninth Revision, Clinical Modification, 2004.

Zero to Three, Diagnostic Classification of Mental Health and Developmental Disorders of Infancy and Early Childhood: Revised Edition (DC: 0-3 R), Washington, DC, 2005.

Maine DC 0-3 Committee, Crosswalk for DC 0-3 to ICD-9 Diagnosis Coding. 2004.

Perez, L. and Newman, M., "Crosswalk" for San Mateo, California, Developmental Approach to Infant Mental Health Diagnosis.

In addition, the Crosswalk developed by Florida was used as a reference.

+ To be used as a guide in Clinical Assessment/Diagnosis